

KURZPROTOKOLL
MS200647-0047

Öffentlicher Titel	Phase II Studie zu M7824 als Zweitlinientherapie bei metastasiertem Gallenwegskrebs
Wissenschaftl. Titel	A Phase II, Multicenter, Open-label Study to Investigate the Clinical Efficacy of M7824 Monotherapy in Participants With Locally Advanced or Metastatic Biliary Tract Cancer Who Fail or are Intolerant to First-line Platinum-Based Chemotherapy
Kurztitel	MS200647-0047
Studienart	multizentrisch, prospektiv, Therapiestudie, offen/unverblindet, einarmig, Pharma-Studie
Studienphase	Phase II
Erkrankung	Verdauung: Gallengangs-/Gallenblasenkrebs (maligne biliäre Tumoren): Zweitlinie oder höher
Einschlusskriterien	- Are participants with histologically or cytologically confirmed locally advanced or metastatic BTC. - Availability of tumor (primary or metastatic) archival material or fresh biopsies (collected within 28 days before first administration) of study intervention is mandatory - Participants with BTC must have failed or be intolerant to 1L systemic platinum-based chemotherapy - Disease must be measurable with at least 1 unidimensionally measurable lesion by RECIST 1.1 - Eastern Cooperative Oncology Group (ECOG) PS of 0 to 1 - Life expectancy ≥ 12 weeks as judged by the Investigator - Adequate hematological function defined by white blood cell (WBC) count $\geq 3 \times 10^9/\text{Litre}$ with absolute neutrophil count (ANC) $\geq 1.5 \times 10^9/\text{Litre}$, lymphocyte count $\geq 0.5 \times 10^9/\text{Litre}$, platelet count $\geq 75 \times 10^9/\text{Litre}$, and hemoglobin (Hgb) ≥ 9 grams/decilitre - Adequate hepatic function defined by a total bilirubin level $\leq 1.5 \times$ upper limit of normal (ULN), an aspartate aminotransferase (AST) level $\leq 2.5 \times$ ULN, and an alanine aminotransferase (ALT) level $\leq 2.5 \times$ ULN. For participants with liver involvement in their tumor, AST $\leq 5.0 \times$ ULN and ALT $\leq 5.0 \times$ ULN is acceptable - Adequate coagulation function defined as prothrombin time (PT) or international normalized ratio (INR) $\leq 1.5 \times$ ULN unless the participant is receiving anticoagulant therapy - Albumin ≥ 3.0 grams/decilitre - Hepatitis B virus (HBV) deoxyribonucleic acid (DNA) positive participants must be treated and on a stable dose of antivirals - Adequate renal function defined by either creatinine $\leq 1.5 \times$ ULN or an estimated creatinine clearance (CCr) > 40 milliliter (mL) per minute (min) according to the Cockcroft-Gault formula or by measure of CCr from 24-hour urine collection - Other protocol defined inclusion criteria could apply
Ausschlusskriterien	- Ampullary cancer is excluded - Significant acute or chronic infections - Active autoimmune disease that might deteriorate when receiving an immunostimulatory agent - Interstitial lung disease or its history - Participants who are not eligible for or have not been treated with 1L systemic chemotherapy - Anticancer treatment within 21 days before the start of study intervention - Concurrent treatment with nonpermitted drugs - Prior participation in a M7824 clinical trial

KURZPROTOKOLL
MS200647-0047

- Prior therapy with other immunotherapy or checkpoint inhibitors, such as anti-PD 1, anti PD L1, anti- cytotoxic T-cell lymphocyte-4 (CTLA-4) antibodies.
- Pregnancy or breast feeding
- Other protocol defined exclusion criteria could apply

Alter 18 Jahre und älter

Sponsor Merck KGaA

Registrierung in anderen Studienregistern EudraCT 2018-003707-19
ClinicalTrials.gov NCT03833661 (primäres Register)