

KURZPROTOKOLL
Napoli

Öffentlicher Titel	Vergleich zweier Erstlinien-Therapieregime für das metastasierte Adenokarzinom der Bauchspeicheldrüse
Wissenschaftl. Titel	An open-label, randomised, multicentre, phase III study of irinotecan liposome injection, oxaliplatin, 5-fluorouracil/leucovorin versus nab-paclitaxel plus gemcitabine in subjects who have not previously received chemotherapy for metastatic adenocarcinoma of the pancreas
Kurztitel	Napoli
Studienart	multizentrisch, prospektiv, Therapiestudie, randomisiert, offen/unverblindet, Register, Pharma-Studie, zweiarmig
Studienphase	Phase III
Erkrankung	Verdauung: Bauchspeicheldrüsenkrebs (Pankreaskarzinom): Erstlinie
Einschlusskriterien	- Histological or cytologically confirmed adenocarcinoma of the pancreas that has not been previously treated in the metastatic setting - Initial diagnosis of metastatic disease must have occurred ≤ 6 weeks prior to screening - Subject has one or more metastatic tumours measurable by computed tomography (CT) scan (or magnetic resonance imaging (MRI), if the subject is allergic to CT contrast media) according to RECIST Version 1.1 criteria - Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1 - Subject has adequate biological parameters as demonstrated by the following blood counts: (a) Absolute neutrophil count (ANC) $\geq 2000/\text{mm}^3$ without the use of hemopoietic growth factors within the last 7 days prior to randomisation (b) Platelet count $\geq 100,000/\text{mm}^3$ (c) Haemoglobin (Hgb) $\geq 9 \text{ g/dL}$ obtained ≤ 14 days prior to randomisation - Adequate hepatic function as evidenced by: (a) Serum total bilirubin $\leq 1.5 \times \text{ULN}$ (biliary drainage is allowed for biliary obstruction), and (b) Aspartate aminotransferase (AST) and alanine aminotransferase (ALT) $\leq 2.5 \times$ upper limit of normal (ULN) ($\leq 5 \times \text{ULN}$ is acceptable if liver metastases are present) - Adequate renal function as evidenced by creatinine clearance $\geq 30 \text{ mL/min}$ - Adequate coagulation studies (obtained ≤ 14 days prior to randomisation) as demonstrated by prothrombin time and partial thromboplastin time within normal limits ($\leq 1.5 \times \text{ULN}$)
Ausschlusskriterien	- Prior treatment of pancreatic cancer in the metastatic setting with surgery, radiotherapy, chemotherapy or investigational therapy - Prior treatment of pancreatic adenocarcinoma with chemotherapy in the adjuvant setting, except those where at least 12 months have elapsed since completion of the last dose and no persistent treatment-related toxicities are present - Subject has only localised advanced disease - Documented serum albumin $< 3 \text{ g/dL}$ - Known history of central nervous system (CNS) metastases - Clinically significant gastrointestinal disorder - History of any second malignancy in the last 2 years - Concurrent illnesses that would be a relative contraindication to trial participation - Use of strong inhibitors or inducers of CYP3A, CYP2C8 and UGT1A1 - Neuroendocrine (carcinoid, islet cell) or acinar pancreatic carcinoma - Known low or absent dihydropyrimidine dehydrogenase (DPD) activity
Alter	18 Jahre und älter

KURZPROTOKOLL
Napoli

Prüfzentren	Krankenhaus Nordwest GmbH (Rekrutierung beendet) Institut für klinisch-onkologische Forschung Steinbacher Hohl 2-26 60488 Frankfurt am Main Markus Knetsch Tel: 069 7601 - 3128 knetsch.markus@khnw.de
Sponsor	Ipsen Pharma
Registrierung in anderen Studienregistern	ClinicalTrials.gov NCT04083235 (primäres Register) EudraCT 2018-003585-14