

## **KURZPROTOKOLL** **PrefMab (MO28457)**

<b>Öffentlicher Titel</b>	Subkutane oder intravenöse Verabreichung von Rituximab bei nicht-vorbehandeltem, CD20+ DLBCL oder follikulärem NHL
<b>Wissenschaftl. Titel</b>	Eine randomisierte, offene, multizentrische Studie zur Bewertung der Patientenpräferenz bei Vergleich der subkutanen und intravenösen Verabreichung von Rituximab bei nicht-vorbehandelten Patienten mit CD20+ diffus großzelligem BZell-Lymphom oder CD20+ follikulärem Non-Hodgkin-Lymphom Grad 1, 2 oder 3a.
<b>Kurztitel</b>	PrefMab (MO28457)
<b>Studienart</b>	multizentrisch, prospektiv, randomisiert, offen/unverblindet, Pharma-Studie, zweiarmig
<b>Studienphase</b>	Phase III
<b>Erkrankung</b>	Blut: Non-Hodgkin-Lymphome (NHL), niedrig-maligne: andere NHL - neu diagnostiziert / de novo
<b>Ziele</b>	<ul style="list-style-type: none"><li>- Proportion of patients indicating an overall preference via Patient Preference Questionnaire (PPQ) for either the subcutaneous (SC) or intravenous (IV) administration of MabThera/Rituxan</li><li>- Safety: Incidence of adverse events</li><li>- Administration time SC vs IV</li><li>- Patient assessed satisfaction SC vs IV: Cancer Therapy Satisfaction Questionnaire (CTSQ)/Rituximab Administration Satisfaction Questionnaire (RASQ)</li><li>- Complete response (CR) rate including complete response unconfirmed (CRu) 4-8 weeks after last dose of induction treatment</li><li>- Event-free survival (EFS)</li><li>- Disease-free survival (DFS)</li><li>- Progression-free survival (PFS)</li><li>- Overall survival (OS)</li><li>- Immunogenicity: Anti-rituximab and anti-human recombinant hyaluronidase [rHuPH20] antibodies, associated rituximab concentration level)</li></ul>
<b>Einschlusskriterien</b>	<ul style="list-style-type: none"><li>- Adult patients, <math>\geq 18</math> and <math>\leq 80</math> years of age</li><li>- Histologically confirmed, previously untreated CD20+ diffuse large B-cell lymphoma (DLBCL) or CD20+ follicular non-Hodgkin's lymphoma (NHL) Grade 1, 2, or 3a, according to WHO classification</li><li>- An International Prognostic Index (IPI) score of 1-4 or IPI score of 0 with bulky disease, defined as one lesion <math>\geq 7.5</math> cm, or Follicular Lymphoma International Prognostic Index (FLIPI; low, intermediate or high risk)</li><li>- At least one bi-dimensionally measurable lesion defined as <math>\geq 1.5</math> cm in its largest dimension on CT scan</li><li>- Eastern Cooperative Oncology Group (ECOG) performance status <math>\leq 3</math></li></ul>
<b>Ausschlusskriterien</b>	<ul style="list-style-type: none"><li>- Transformed lymphoma or follicular lymphoma IIIB</li><li>- Primary central nervous system (CNS) lymphoma, blastic variant of mantle-cell lymphoma, histologic evidence of transformation to Burkitt lymphoma, primary mediastinal DLBCL, primary effusion lymphoma, primary cutaneous DLBCL, or primary DLBCL of the testis</li><li>- History of other malignancy that could affect compliance with the protocol or interpretation of the results; this includes a malignancy that has been treated but not with curative intent, unless the malignancy has been in remission for <math>\geq 5</math> years prior to enrolment; patients with a history of curatively treated basal or squamous cell carcinoma or melanoma of the skin or in situ carcinoma of the cervix are eligible</li><li>- Prior therapy for DLBCL or NHL, with the exception of nodal biopsy or local irradiation</li></ul>

**KURZPROTOKOLL**  
**PrefMab (MO28457)**

- Prior treatment with cytotoxic drugs (with the exclusion of methotrexate for CNS prophylaxis in DLBCL) or rituximab for another condition, or prior use of an anti-CD20 drug
- Prior use of monoclonal antibody within 3 months prior to randomization
- Chemotherapy or other investigational therapy within 28 days prior to randomization
- Ongoing corticosteroid use > 30 mg/day prednisolone or equivalent
- Inadequate renal, hematologic or hepatic function
- Active and/or severe infection or any major episode of infection within 4 weeks prior to randomization
- Active hepatitis B virus or active hepatitis C virus infection
- History of human immunodeficiency (HIV) seropositive status
- A positive pregnancy test in women of childbearing potential
- Life expectancy of less than 6 months

<b>Alter</b>	18 - 80 Jahre
<b>Molekularer Marker</b>	CD20
<b>Fallzahl</b>	900
<b>Sponsor</b>	Hoffmann-La Roche (Hauptsponsor)
<b>Förderer</b>	Hoffmann-La Roche
<b>Registrierung in anderen Studienregistern</b>	ClinicalTrials.gov NCT01724021 EudraCT 2012-003230-17