

KURZPROTOKOLL ARAMIS

Öffentlicher Titel	Phase-III-Studie zur Wirksamkeit und Sicherheit von ODM-201 bei männlichen Hochrisikopatienten mit nicht metastasiertem, kastrationresistentem Prostatakrebs
Wissenschaftl. Titel	Multizentrische, randomisierte, doppelblinde, placebokontrollierte Phase-III-Studie zur Wirksamkeit und Sicherheit von ODM-201 bei männlichen Hochrisikopatienten mit nicht metastasiertem, kastrationresistentem Prostatakrebs
Kurztitel	ARAMIS
Studienart	multizentrisch, randomisiert, Pharma-Studie, doppelblind, zweiarmig
Studienphase	Phase III
Erkrankung	Geschlechtsorgane: Krebserkrankungen der männlichen Geschlechtsorgane: Prostatakrebs - Erstlinie
Einschlusskriterien	- Castration-resistant prostate cancer (CRPC) with castrate level of serum testosterone. - Prostate-specific antigen doubling time of 10 months and PSA > 2ng/ml. - Eastern Cooperative Oncology Group (ECOG) performance status of 0-1. - Blood counts at screening: haemoglobin 9.0 g/dl, absolute neutrophil count 1500/μl, platelet count 100,000/μl. - Screening values of serum alanine aminotransferase (ALT) and/or aspartate transaminase (AST) 2.5 x upper limit of normal (ULN), total bilirubin 1.5 x ULN, creatinine 2.0 x ULN. - Sexually active patients, unless surgically sterile, must agree to use condoms as an effective barrier method and refrain from sperm donation during the study treatment and for 3 months after the end of the study treatment.
Ausschlusskriterien	- History of metastatic disease or presence of detectable metastases. - Acute toxicities of prior treatments and procedures not resolved to grade 1 or baseline before randomisation. - Prior treatment with: second generation androgen receptor (AR) inhibitors, other investigational AR inhibitors, or CYP17 enzyme inhibitor. - Use of estrogens, 5- reductase inhibitors or AR inhibitors. - Prior chemotherapy or immunotherapy for prostate cancer. - Use of systemic corticosteroid. - Radiation therapy within 12 weeks before randomisation. - Severe or uncontrolled concurrent disease, infection or co-morbidity. - Treatment with bisphosphonate or denosumab within 12 weeks before randomisation. - Known hypersensitivity to the study treatment or any of its ingredients. - Major surgery within 28 days before randomisation. - Any of the following within 6 months before randomisation: stroke, myocardial infarction, severe/unstable angina pectoris, coronary/peripheral artery bypass graft; congestive heart failure New York Heart Association (NYHA) Class III or IV. - Uncontrolled hypertension. - Prior malignancy. - Gastrointestinal disorder or procedure which expects to interfere significantly with absorption of study treatment. - Active viral hepatitis, active human immunodeficiency virus (HIV) or chronic liver disease. - Any condition that in the opinion of the investigator would impair the patients' ability to comply with the study procedures.
Alter	18 Jahre und älter

KURZPROTOKOLL
ARAMIS

Fallzahl	1500
Prüfzentren	Universitätsmedizin Frankfurt (Rekrutierung beendet) Zentrum für Chirurgie, Klinik für Urologie Theodor-Stern-Kai 7 60590 Frankfurt am Main Dr. med. Severine Banek Tel: 069 6301-80072 Fax: 069 6301-84029 severine.banek@unimedizin-ffm.de
Sponsor	Bayer Healthcare
Registrierung in anderen Studienregistern	ClinicalTrials.gov NCT02200614 EudraCT 2013-003820-36