

KURZPROTOKOLL
2215-CL-0201

Öffentlicher Titel	Gilteritinib und Azacitidin bei nicht-intensiv behandelbarer de novo AML mit FLT3 Mutation
Wissenschaftl. Titel	A Phase 2/3 Multicenter, Open-label, 3-arm, 2-stage Randomized Study of ASP2215 (Gilteritinib), Combination of ASP2215 Plus Azacitidine and Azacitidine Alone in the Treatment of Newly Diagnosed Acute Myeloid Leukemia with FLT3 Mutation in Patients Not Eligible for Intensive Induction Chemotherapy
Kurztitel	2215-CL-0201
Studienart	prospektiv, Therapiestudie, randomisiert, offen/unverblindet, dreiarmlig
Studienphase	Phase II/III
Erkrankung	Blut: Akute myeloische Leukämie (AML): Neu diagnostiziert / de novo
Einschlusskriterien	<ul style="list-style-type: none">- Subject is considered an adult according to local regulation at the time of obtaining informed consent.- Subject has a diagnosis of previously-untreated AML according to World Health Organization (WHO) classification [Swerdlow et al, 2008] as determined by pathology review at the treating institution.- Subject is positive for FLT3 mutation (internal tandem duplication [ITD] or tyrosine kinase domain [TKD] [D835/I836] mutation) in bone marrow or whole blood as determined by central laboratory. Note: Only applicable to the randomization portion.- Subject has received a cumulative anthracycline dose above 400 mg/m2 of doxorubicin (or cumulative maximum dose of another anthracycline).- Subject must meet the following criteria as indicated on the clinical laboratory tests: Serum AST and ALT $\leq 2.5 \times$ Institutional upper limit of normal (ULN); (b) Serum total bilirubin $\leq 1.5 \times$ Institutional ULN; (c) Serum potassium \geq Institutional lower limit of normal (LLN); (d) Serum magnesium Institutional LLN- Subject is suitable for oral administration of study drug.- Female subject must either: (a) Be of nonchildbearing potential: (1) Postmenopausal (defined as at least 1 year without any menses) prior to screening, or (2) Documented surgically sterile or status posthysterectomy (at least 1 month prior to screening); (b) Or, if of childbearing potential, (1) Agree not to try to become pregnant during the study and for 180 days after the final study drug administration (2) And have a negative urine or serum pregnancy test at screening (3) And, if heterosexually active, agree to consistently use 2 forms of effective contraception per locally accepted standards, 1 of which must be a barrier method, starting at screening and throughout the study period and for 180 days after the final study drug administration.- Female subject must agree not to breastfeed starting at screening and throughout the study period, and for 60 days after the final study drug administration.- Female subject must not donate ova starting at screening and throughout the study period, and for 180 days after the final study drug administration.- Male subject and their female partners who are of childbearing potential must be using 2 forms of effective contraception per locally accepted standards, 1 of which must be a barrier method, starting at screening and continue throughout the study period, and for 120 days after the final study drug administration.- Male subject must not donate sperm starting at screening and throughout the study period and for 120 days after the final study drug administration.- Subject agrees not to participate in another interventional study while on treatment.
Ausschlusskriterien	<ul style="list-style-type: none">- Subject was diagnosed as acute promyelocytic leukemia (APL).- Subject has BCR-ABL-positive leukemia (chronic myelogenous leukemia in blast crisis).

KURZPROTOKOLL
2215-CL-0201

- Subject has received previous therapy for AML, with the exception of the following: (a) Emergency leukapheresis; (b) Hydroxyurea for ≤ 14 days; (c) Preemptive treatment with retinoic acid prior to exclusion of APL ≤ 7 days; (d) Growth factor or cytokine support; (e) Steroids for the treatment of hypersensitivity or transfusion reactions, nausea/vomiting or pain
- Subject has clinically active central nervous system leukemia.
- Subject has been diagnosed with another malignancy that requires concurrent treatment or hepatic malignancy regardless of need for treatment.
- Subject has clinically significant coagulation abnormality unless secondary to AML.
- Subject has had major surgery within 4 weeks prior to the first study dose.
- Subject has radiation therapy within 4 weeks prior to the first study dose.
- Subject requires treatment with concomitant drugs that are strong inducers of cytochrome P450 CYP3A.
- Subject requires treatment with concomitant drugs that are strong inhibitors or inducers of P-gp with the exception of drugs that are considered absolutely essential for the care of the subject.
- Subject requires treatment with concomitant drugs that target serotonin 5HT1R or 5HT2BR or sigma nonspecific receptor with the exception of drugs that are considered absolutely essential for the care of the subject.
- Subject has congestive heart failure classified as New York Heart Association Class IV.
- Subject with mean Fridericia-corrected QT interval (QTcF) > 450 ms at screening based on central reading.
- Subject with a history of Long QT Syndrome at screening.
- Subject has known pulmonary disease with diffusion capacity of lung for carbon monoxide (DLCO) $\leq 65\%$, forced expiratory volume in the first second (FEV1) $\leq 65\%$, dyspnea at rest or requiring oxygen or any pleural neoplasm (Transient use of supplemental oxygen is allowed.)
- Subject has an active uncontrolled infection. If an infection is present, the patient must be receiving definitive therapy and have no signs of progressing infection. Progressing infection is defined as hemodynamic instability attributable to sepsis or new symptoms, worsening physical signs or radiographic findings attributable to infection. Persisting fever without other signs or symptoms will not be interpreted as progressing infection.
- Subject is known to have human immunodeficiency virus infection.
- Subject has active hepatitis B or C or other active hepatic disorder.
- Subject has any condition which makes the subject unsuitable for study participation, including any contraindications of azacitidine.

Alter

18 - 70 Jahre

Molekularer Marker

FLT3

Prüfzentren

Innere Medizin 2 (Geschlossen)
Hämatologie / Medizinische Onkologie
Theodor-Stern-Kai 7
60590 Frankfurt am Main
Silvia Koss
Tel: 069 6301-80429
Fax: 069 6301-83655
silvia.koss@unimedizin-ffm.de

Sponsor

Astellas Pharma

KURZPROTOKOLL
2215-CL-0201

**Registrierung in anderen
Studienregistern**

ClinicalTrials.gov NCT02752035
EudraCT 2015-001790-41

Links

[Studiendokumente zum Download \(roXtra\)](#)